



Review

Gaze sensitivity: function and mechanisms from sensory and cognitive perspectives



Gabrielle L. Davidson^{a,*}, Shannon Butler^b, Esteban Fernández-Juricic^b, Alex Thornton^{a,c}, Nicola S. Clayton^a

^a Department of Psychology, University of Cambridge, Cambridge, U.K.

^b Department of Biological Sciences, Purdue University, West Lafayette, IN, U.S.A.

^c Centre for Ecology and Conservation–Biosciences, University of Exeter, Penryn, U.K.

ARTICLE INFO

Article history:

Received 27 January 2013

Initial acceptance 27 February 2013

Final acceptance 30 September 2013

Available online 21 November 2013

MS. number: 13-00081R

Keywords:

cognition
gaze aversion
gaze following
gaze sensitivity
retina
visual field
visual fixation

Sensitivity to the gaze of other individuals has long been a primary focus in sociocognitive research on humans and other animals. Information about where others are looking may often be of adaptive value in social interactions and predator avoidance, but studies across a range of taxa indicate there are substantial differences in the extent to which animals obtain and use information about other individuals' gaze direction. As the literature expands, it is becoming increasingly difficult to make comparisons across taxa as experiments adopt and adjust different methodologies to account for differences between species in their socioecology, sensory systems and possibly also their underlying cognitive mechanisms. Furthermore, as more species are found to exhibit gaze sensitivity, more terminology arises to describe the behaviours. To clarify the field, we propose a restricted nomenclature that defines gaze sensitivity in terms of observable behaviour, independent of the underlying mechanisms. This is particularly useful in nonhuman animal studies where cognitive interpretations are ambiguous. We then describe how socioecological factors may influence whether species will attend to gaze cues, and suggest links between ultimate factors and proximate mechanisms such as cognition and perception. In particular, we argue that variation in sensory systems, such as retinal specializations and the position of the eyes, will determine whether gaze cues (e.g. head movement) are perceivable during visual fixation. We end by making methodological recommendations on how to apply these variations in socioecology and visual systems to advance the field of gaze research.

© 2013 The Association for the Study of Animal Behaviour. Published by Elsevier Ltd. All rights reserved.

Attending to where others are looking may offer important information about the location of food and predators, as well as social relationships between conspecifics. Humans show gaze sensitivity in many contexts: we can accurately follow where others are looking in space (e.g. Bock, Dicke, & Their, 2008), and appreciate that others may have different fields of view or perspectives. We use our own gaze as a form of communication to inform or mislead others, and use the gaze of others to interpret their mental states (e.g. Teufel, Alexis, Clayton, & Davis, 2010).

A number of other species including mammals, birds and reptiles have also been reported to show sensitivity to gaze. Sensitivity to gaze can result in many different responses, such as avoiding gaze because it is associated with the approach of a predator, or co-orienting with another's gaze to spot objects of interest. Behavioural and sensory ecologists have sought to determine the socioecological contexts in which gaze sensitivity occurs, and to

identify features of cues that are most important for eliciting gaze sensitivity responses (e.g. Burger, Gochfeld, & Murray, 1991; Carter, Lyons, Cole, & Goldsmith, 2008; Hampton, 1994; Watve et al., 2002). Numerous experimental paradigms have also been developed to test whether these responses are simply reflexive, and therefore bound to one stimulus in one context, or whether they involve further information processing (e.g. von Bayern & Emery, 2009a; Bugnyar, Stowe, & Heinrich, 2004; Loretto, Schloegl, & Bugnyar, 2010). The study of this information processing has been of great interest to cognitive psychologists (e.g. Call, Hare, & Tomasello, 1998; Povinelli & Eddy, 1996). Many tasks have been designed to identify the cognitive mechanisms by which information from another's direction of attention is processed, and whether these mechanisms allow subjects to apply gaze information flexibly in different contexts, and/or through different behavioural responses. As a result, a plethora of experimental paradigms have been developed to address gaze behaviours in a multitude of different species and contexts.

The aim of this review is two-fold. The first goal is to present a standardized set of nomenclature that brings together all aspects of

* Correspondence: G. Davidson, Department of Psychology, University of Cambridge, Downing Street, Cambridge CB2 3EB, U.K.

E-mail address: gd339@cam.ac.uk (G. L. Davidson).

gaze research (gaze preference, gaze following and gaze aversion), and defines these behaviours independently from cognitive mechanisms. We hope that this nomenclature brings clarity to the gaze sensitivity literature, and facilitates a bridge between various aspects of gaze research across many disciplines. The second goal is to illustrate how socioecological pressures and proximate anatomical, sensory and cognitive factors can influence the occurrence of gaze sensitivity across taxa. These factors can vary substantially between species, and as the breadth of species studied in gaze contexts increases, it is important to consider this variability when interpreting results, designing gaze sensitivity experiments, and choosing appropriate study species.

DEFINING GAZE BEHAVIOURS

A number of different gaze behaviours have been described in the literature and, as a result, this has brought a sense of confusion because many species are studied in different contexts and some definitions carry with them an assumption of the underlying cognitive processing. For example, an animal may orient their gaze with another individual because they understand the referential nature of looking, i.e. that another individual can see something. Alternatively, an animal may orient their gaze in response to another individual's gaze because having done so in the past resulted in seeing an interesting object. These two scenarios are guided by different processes (discussed in more detail below), but elicit the same observable behaviour. It is therefore useful, particularly in nonhuman research where mental processes are difficult to ascertain, to describe gaze behaviours purely in terms of the observable behaviour. The terminology used should be independent from any

assumptions about the cognitive processes, be it a reflexive response or one that requires further information processing (see [Thornton & Raihani, 2008](#) and [Thornton & McAuliffe, 2012](#) for similar arguments concerning the definition of teaching). This is particularly useful in a field in which multiple disciplines study gaze sensitivity. For those studying underlying cognition, experimental paradigms can be applied specifically to test information processing mechanisms underlying gaze behaviours (as defined below). Here we present nomenclature derived from the literature which we propose be restricted to the following definitions.

Gaze Sensitivity

We propose that all instances whereby an individual attends to gaze stimuli should be classed under the umbrella category of gaze sensitivity. Sensitivity to gaze is a prerequisite for all gaze response behaviours defined below. Whether an individual is sensitive to the gaze of others may be dependent on a number of factors which are discussed throughout this review, including sociality, ecology, cognition and visual architecture. Gaze sensitivity is also dependent upon the gaze cues available.

Gaze Cues

Gaze sensitivity and the resulting gaze behaviours are reliant on an observable gaze cue. Gaze cues include the presence or orientation of the eyes or head, and may be presented as static or moving stimuli. The head and the eyes can be presented in alignment (congruent), or in opposing directions (incongruent), and may also be relative to body positioning. Direct gaze ([Fig. 1a](#)) refers to an individual's gaze




Figure 1. Gaze cues and behaviours. Arrows depict direction of gaze. (a) Direct gaze (single arrow) and mutual gaze (double arrow); (b) direct gaze cue resulting in averted gaze response; (c) gaze following; (d) joint attention; (e) geometric gaze.

directed towards another individual, whereas averted gaze refers to an individual's gaze directed away from another individual. Direct and averted gaze can refer to the cues given, but may also be described as gaze responses (e.g. an individual averts its gaze in response to direct gaze, Fig. 1b). In some cases gaze cues and responses occur between conspecifics or between heterospecifics (e.g. human demonstrator presenting cues to an animal subject, or a predator giving away cues to an animal subject). We now describe gaze behaviours typically observed in response to gaze cues.

Gaze Responses

Gaze sensitivity can result in a number of different gaze responses. These include gaze preference, gaze aversion and gaze following responses. Gaze preference refers to an individual's preference for looking at a particular gaze cue. For example, an individual may spend more time looking at another individual that is looking towards them (direct gaze) than one that is looking away from them (averted gaze), or vice versa. Gaze aversion refers to aversive behaviour in response to the presence of gaze cues, for example an individual moving away from another individual that is looking towards it. Gaze following refers to the act of orienting one's gaze in the direction of another's gaze (Fig. 1c). For example, one individual moves its head to look to the side, and in response, a second individual moves its head in a similar direction. Gaze preference, gaze aversion and gaze following can be further subdivided within these responses (Fig. 2).

Gaze Preference

Gaze preference responses refer to looking behaviour from the subject. When presented with a choice between demonstrators exhibiting different gaze cues, an individual may spend more time looking at an individual showing a preferred gaze cue. Gaze preferences may also result in shorter latencies for spotting individuals in a crowd displaying particular gaze cues. For instance, Tomonaga and Imura (2010) showed that when an adult chimpanzee, *Pan troglodytes*, was presented with a screen of many human faces, the subject was faster at detecting a face with direct eye gaze than a face with averted eye gaze. When presented with only one demonstrator, gaze preference may be directed to a specific area of

the face such as the eyes rather than the head in general. The demonstrator and the subject may engage in mutual gaze, where both individuals look at one another (Fig. 1a).

Gaze Aversion

In gaze aversion, the possible behaviours may be reliant on the context in which the gaze cues are presented. A sudden appearance or approach of gaze cues can elicit aversive escape responses, generally associated with antipredator responses such as fleeing, crouching or tonic immobility. Similar responses such as fleeing or looking away may also occur between conspecifics, for instance between individual territory holders, or within dominance hierarchies. Gaze aversion can also include behaviours in which an animal is approaching, as opposed to when it is moving away. We refer to aversive approach if a gaze cue is directed towards a desired object such as food, and the subject alters its behaviour by delaying its approach, or approaching only when the gaze cue is averted or hidden.

Gaze Following

In gaze following, individuals may orient their gaze in the same direction, but this does not imply they are necessarily looking at the same thing. In its simplest form, gaze following refers to the co-orientation of gaze with another towards a similar point in space (Emery, Lorincz, Perrett, Oram, & Baker, 1997). Following Emery et al. (1997), Emery (2000), we distinguish gaze following from joint attention. In the latter, an individual not only orients its gaze in the same direction of another's, but as a result, both individuals' gazes are directed towards the same object (Fig. 1d). This does not suggest that those engaging in joint attention must appreciate the visual attention of others. Further testing would be necessary to pinpoint the cognitive mechanisms (see below). As well as orienting one's gaze with another, an individual may need to reposition itself to be in the same line of sight as the demonstrator. In geometric gaze, an individual repositions itself around a barrier to follow the gaze of another individual (Fig. 1e). Geometric gaze may result in joint attention if both individuals subsequently gaze at the same thing behind the barrier.

This terminology serves to bring together all aspects of gaze research. Behaviours such as gaze aversion and gaze following are




Figure 2. Diagram depicting proposed gaze nomenclature. Gaze sensitivity is reliant on the gaze cues available. Sensitivity to gaze cues will result in gaze behaviours that are described within the categories of gaze preference, gaze aversion and gaze following.

often studied separately (but see von Bayern & Emery, 2009a), yet are interrelated in that they rely on/are based on animals' responses to gaze cues. Therefore it is useful to use the term gaze sensitivity when discussing responses to gaze cues in a broad context, and to use the additional behavioural definitions when discussing more specific responses to gaze. Our nomenclature describes the basic components of gaze tasks in terms of behaviours without assumptions about unobservable underlying mechanisms. Once behavioural responses have been observed and categorized, tests can be designed to tease apart the underlying processes that guide these behaviours (cf. Thornton & McAuliffe, 2012; Thornton & Raihani, 2008). For instance, do individuals consider where another individual's direction of attention is focused? Might they recognize that another individual's line of sight may be different from their own? Can they use another individual's gaze to infer that individual's intention towards an object? Are individuals able to use gaze flexibly by applying different behavioural responses or cognitive mechanisms across different contexts (e.g. to detect predator gaze, to follow conspecific gaze to find food and to find predators), or are they bound to one particular response in one particular context? An individual's gaze response may also be dependent upon the availability of gaze cues and their characteristics. For instance, some species may be more sensitive to head direction because they move their head more than their eyes when scanning for or fixating on objects. Alternatively, some species may gain more information from the eyes than the head. Species' differences in available gaze cues (e.g. rate and/or orientation of eye or head movement) are highly dependent upon the configuration of the animal's visual system.

Carefully designed experiments allow us (1) to determine how the sensory system of a given species gathers gaze information and (2) to establish the cognitive requirements for different gaze behaviours. These proximate mechanisms may help to explain why we see variation in gaze following and gaze aversion behaviours across species. It is equally important to consider ultimate mechanisms, namely socioecological factors that will determine whether attending to gaze cues is beneficial to the observer. Variability in socioecological pressures may in fact drive species to process gaze cues such that they can be applied across various contexts. Because this may also be a function of the species' underlying cognition and sensory system, we expect proximate and ultimate mechanisms of gaze sensitivity to be linked, and therefore should be studied in concert.

SOCIOECOLOGY AND CUE INFORMATION

Consideration of socioecological factors is essential to understand the selection pressures driving the evolution of different forms of gaze sensitivity behaviours. Moreover, socioecological considerations also provide critical information into the proximate basis of gaze sensitivity. We expect sensitivity to gaze to occur only if cues are discernible and provide useful information on which the observer can act. Therefore there is often interplay between socioecological contexts and the features of the gaze cues available. For instance, predator detection may be dependent on the salience of the predator's eyes, or the prey's capacity to perceive the gaze cues of a heterospecific. There may be a selection pressure for predators to evolve less conspicuous eyes, or to evolve visual configurations that are different from their prey species, making detection of predator gaze more difficult. Similarly, experiments testing for gaze sensitivity often differ in their use of heterospecific (human, predator) or conspecific demonstrators, which may affect whether the subject is motivated to attend to the demonstrator (Bräuer, Call, & Tomasello, 2005; Bugnyar et al., 2004; Emery et al., 1997; Tomasello, Call, & Hare, 1998). Therefore socioecology can give insight into the underlying mechanisms that facilitate the occurrence of gaze behaviours.

Gaze Cues from Predators

A predator's gaze may give prey species accurate information about the necessity of escape. By accurately assessing where a predator is looking, species may ultimately benefit from increased foraging opportunities (Carter et al., 2008) or more frequent nest visits (Watve et al., 2002). Risk perception may be influenced by the properties of the gaze cue provided by the predator, such as the positioning of the head or eyes, and the colour, shape and size of the eyes (Burger et al., 1991; Coss, 1979; Jones, 1980; Scaife, 1976a). Enhancing or presenting contradictory cues can help experimenters isolate important stimuli for aversive escape responses. House sparrows, *Passer domesticus*, fly away most when a human model is facing towards them, but attend only to head orientation rather than eye orientation (Hampton, 1994). Black iguanas, *Ctenosaura similis*, for example, move away sooner when a human face is visible, rather than covered with hair during approach (Burger & Gochfeld, 1993). Similar increases in vigilant behaviours are found when the eyes are made to appear larger (Burger et al., 1991). Two eye-like stimuli horizontally placed side-by-side elicit the most fearful responses in jewel fish, *Hemichromis bimaculatus* (Coss, 1979), while in domestic chicks, *Gallus gallus*, the pairing of an iris with a pupil shape (i.e. having the features of an eye) increases aversive responses (e.g. freezing, distress calls, number of approaches; Jones, 1980) in comparison to other spot arrangements such as no iris or only one eye. However, when testing a small passerine's preference for invertebrates, there is evidence to suggest that any conspicuous shape, such as a square or triangle on the wings of moths, may be as effective as eye-shaped spots in deterring predation (Stevens et al., 2007).

Gaze cues that elicit fearful responses may also be important if an animal must approach an object or area where a dangerous agent (e.g. unfamiliar human or predator) is gazing. The conflict paradigm tests whether the subject attends to the orientation of the experimenter's head or eyes by measuring its latency to approach a desired item such as food. If subjects refrain from approaching the food for some time this suggests they are fearful of the experimenter and potentially regard them as a threat. If the subject is attending to gaze, the latency to approach is expected to be longest when the experimenter is looking towards the object (e.g. von Bayern & Emery, 2009a; Carter et al., 2008). This paradigm has mainly been tested on birds, perhaps because of their vigilant, flighty behaviour in the presence of a dangerous agent (typically a human experimenter) alongside their willingness to approach food. Green bee-eaters, *Merops orientalis*, approach their nest sites less (Watve et al., 2002) and starlings, *Sturnus vulgaris*, (Carter et al., 2008) are less likely to approach food sources when a human experimenter is looking. Jackdaws, *Corvus monedula*, show similar responses to starlings, but only if the experimenter is unfamiliar (von Bayern & Emery, 2009a). Starlings and jackdaws attend specifically to eye orientation of a different species, not just head orientation.

Assessing a predator's gaze is likely to be constrained by distance effects, which reduce visual contrast and thus limit the ability to perceive subtle cues (Fernández-Juricic & Kowalski, 2011) such as gaze. Individuals may need to get closer to a predator to determine its gaze direction, which could increase predation risk. Consequently, we would expect that sensitivity to predator eye gaze would be more likely in species with high visual acuity (i.e. large eye size relative to body mass, presence of a fovea) as they would be able to resolve at further distances variations in the predator's behaviour without incurring too much risk.

The studies cited above examined differential responses to head or eye movement between heterospecifics (i.e. between the subject and the predator or unfamiliar human), but there are also instances of aversive responses between conspecifics. Chimpanzees (Hare, Call,

Agnetta, & Tomasello, 2000) and common marmosets, *Callithrix jacchus* (Burkart & Heschl, 2007) prefer to approach food that a dominant individual cannot see. However, the gaze cues available between conspecifics may not reflect the cues available between heterospecifics (i.e. prey and predator). For instance, chimpanzees and common marmosets may be less sensitive to information from the eyes of conspecifics than humans are, perhaps because many primates have morphological features thought to conceal gaze direction (i.e. dark or no exposed sclera; Kobayashi & Kohshima, 1997, 2001; Tomasello, Hare, Lehmann, & Call, 2007). Characterizing the features of a species' sensory system is necessary in determining what gaze cues are available between conspecifics and heterospecifics.

Gaze Cues from Group Members in Predator Detection

Information about potential predation risk may be gained not only from the predator but also from the gaze of other group members. Many theoretical models of predator avoidance in single- and mixed-species groups assume that collective detection is behind the transfer of information between individuals about potential predator attacks (e.g. Lima, 1987). One possibility is that this transfer of information may also occur through gaze following. When animals are further away in a group, they orient their heads more towards groupmates, possibly to gather information (Fernández-Jurícic, Smith, & Kacelnik, 2005). Studies on primates (Tomasello et al., 1998), birds (Kehmeier, Schloegl, Scheiber, & Weiß, 2011; Loretto et al., 2010), goats, *Capra hircus* (Kaminski, Riedel, Call, & Tomasello, 2005) and the red-footed tortoise, *Chelonoidis carbonaria* (Wilkinson, Mandl, Bugnyar, & Huber, 2010) show that individuals follow the gaze of conspecifics looking up, suggesting they attend to conspecifics as a means to detect aerial predators. Following look-ups of group members may be particularly important for animals that forage by grazing or pecking on the ground. Direction of attention would be divided between food sources (on the ground), predators (e.g. on the horizon or in the sky) and possibly conspecific behaviours (e.g. vigilant look-ups). The necessity of relying on conspecific gaze to detect predators and the availability of information from group members will depend on the animal's visual field. Species with larger visual fields may be able to spot predators when their head is down, while other species may need to look up in order to scan for predators (Fernández-Jurícic, Erichsen, & Kacelnik, 2004).

We have described two aspects of gaze sensitivity that may function in predator avoidance. Both gaze aversion and gaze following behaviours have been reported across a broad spectrum of taxonomic groups, from primates to turtles, and it has been suggested that gaze sensitivity might have been present in a common vertebrate ancestor (Fitch, Huber, & Bugnyar, 2010). However, we note that few studies have yet to investigate predator gaze sensitivity (but see Stevens et al., 2007), for instance, whether predators prefer to approach prey with averted gaze rather than direct gaze. It also remains unclear whether within-species gaze sensitivity is a prerequisite to between-species gaze sensitivity, and whether gaze aversion is a prerequisite to gaze following, or if they are all independent processes. Studies that consider the visual architecture of a species, and apply a variety of paradigms to the same study species using conspecifics and heterospecifics will help decipher whether gaze preference, gaze aversion and gaze following involve the same proximate mechanisms, and whether they evolved dependently or independently.

Social Contexts of Gaze Following

Individuals may gain information from group members by co-orienting their gaze with others, and many species, including all

great apes (Bräuer et al., 2005), rhesus macaques, *Macaca mulatta* (Emery et al., 1997), rooks, *Corvus frugilegus* (Schmidt, Scheid, Kotschal, Bugnyar, & Schloegl, 2011) and ravens, *Corvus corax* (Bugnyar et al., 2004), have been reported to adjust their head direction to match that of a demonstrator. To establish whether individuals are in fact taking into account another individual's visual perspective (as opposed to, for example, behavioural coordination of head movements) experimenters have used the geometric gaze task. In this task, subjects must reorient themselves so they are in line with another individual's field of view, rather than stopping at the first object in sight (i.e. the barrier; Povinelli & Eddy, 1996; Tomasello, Hare, & Agnetta, 1999). One interpretation is that geometric gaze may be useful for species that conceal information or attempt to obtain hidden information from conspecifics. Geometric gaze has been demonstrated in all five great apes (Bräuer et al., 2005; Tomasello et al., 1999), in spider monkeys, *Ateles geoffroyi*, and capuchin monkeys, *Cebus apella* (Amici, Aureli, Visalberghi, & Call, 2009), domestic dogs, *Canis lupus familiaris* (Bräuer, Call, & Tomasello, 2004) and in ravens (Bugnyar et al., 2004). In contrast, northern bald ibises, *Geronticus eremita* (Loretto et al., 2010) and gibbons, *Hylobates* spp. and *Symphalangus syndactylus* (Liebal & Kaminski, 2012) did not gaze behind barriers, indicating that this behaviour is not as widespread as basic gaze following, nor can it be explained by phylogeny as lower apes do not show geometric gaze, while some monkeys do (however, see sensory caveats with regard to gaze sensitivity below). Primates living in competitive social groups may conceal information, for instance by withholding food calls (e.g. Hauser, 1992) or concealing extrapair copulations (le Roux, Snyder-Mackler, Roberts, Beehner, & Bergman, 2013). Gibbons live in small monogamous family groups which may reduce the necessity to conceal actions by group members, although occasional extrapair copulations have been reported (Sommer & Reichard, 2000). The importance of concealment of visual information could be tested by studying geometric gaze in primate species in which same-species individuals may vary in their social dynamics (e.g. male bachelor groups versus family groups). Other lineages known to conceal information from conspecifics include the corvids; therefore, geometric gaze following may be particularly relevant when engaging in caching and pilfering behaviours (Bugnyar et al., 2004; Schloegl, Kotschal, & Bugnyar, 2007).

Some food-caching corvids have been reported to withhold visual and auditory information from potential pilferers (e.g. Bugnyar & Kotschal, 2002; Dally, Emery, & Clayton, 2005; Shaw & Clayton 2012, 2013; Stulp, Emery, Verhulst, & Clayton, 2009), or gain visual information from cachers by preferentially watching conspecifics that are caching, as opposed to conspecifics engaged in noncaching behaviours (Grodzinski, Watanabe, & Clayton, 2012). In a caching paradigm with ravens, a subject observed a human cache two items, while a demonstrator raven was visible to the subject during both caching events, yet had visual access to only one caching event owing to the positioning of a curtain. When given the opportunity to pilfer before their competitor (the demonstrator), subjects preferred to retrieve the food item that was cached when the competitor had visual access, and had no preference when the competitor had no visual access (Bugnyar, 2010). Although these studies did not test behaviour specifically in response to gaze cues, they highlight the importance of a competitor's line of sight during caching and pilfering. Determining whether ravens use gaze cues to find food has been explored explicitly using the object choice task (Schloegl, Kotschal, & Bugnyar, 2008a, 2008b).

In the object choice task, a subject must find food hidden in one of two locations, often under cups or behind barriers. A demonstrator looks in the direction of where the food is hidden, and subjects may attend to the direction of the experimenter or conspecific demonstrator's gaze to determine where food is hidden

(e.g. Call, Agnetta, & Tomasello, 2000; Schloegl et al., 2008a). Ravens were unsuccessful in the object choice paradigm regardless of whether the demonstrator was a conspecific or a human (Schloegl et al., 2008a). Rhesus macaques and capuchin monkeys were also unsuccessful in the object choice task when presented with human gaze cues, although capuchins and some macaques choose above chance when given pointing cues (Anderson, Montant, & Schmitt, 1996; Anderson, Sallaberry, & Barbier, 1995). Chimpanzees also typically perform poorly, perhaps because the experiment is presented in a cooperative framework (Hare & Tomasello, 2004). Chimpanzees are accustomed to frequent competition with group members for access to food (e.g. Hare, Call, & Tomasello, 2006; Hauser, Teixidor, Fields, & Flaherty, 1993), and may not use altruistic, communicative gaze cues. Modifications to the object choice task can often influence success rates, for instance ensuring the demonstrator, rather than the cups, is the main target of the subject's attention. In a meta-analysis of existing object choice tasks using gaze cues (and pointing gestures), success rates were higher if the subject was kept at a distance, or restrained until the cues had been presented for a given period of time before allowing the subject to make a choice (Mulcahy & Hedge, 2012). Therefore performance levels may be attributed to methodological issues involving the salience of the cue or the configuration of the sensory system (see below), rather than a species' cognitive capacity to pass the object choice task.

The object choice task first requires joint attention behaviour as the subject must attend to the same object as the experimenter. Looking at the same cup as the demonstrator (i.e. joint attention) may be achieved by gaze following, and then by visually fixating on the nearest object in sight. Alternatively, looking at the same cup as the demonstrator may be achieved through shared attention, a mechanism involving awareness that one shares attention with another individual towards the same object (Baron-Cohen, 1994; Emery, 2000). In addition to fixating on a particular cup, subjects tested in the object choice task must also use this information subsequently to choose a cup to obtain the hidden reward. A number of researchers have proposed that social interactions involving shared attention may also involve joint intention, a mechanism allowing others to be perceived as intentional agents, and enabling one to form a cognitive representation of one's own intention as well as another individual's intention towards the same object or goal (Tomasello & Carpenter 2005; Tomasello, Carpenter, Call, Behne, & Moll, 2005). Together, shared attention and joint intention can enable shared intentionality in which individuals engage in collaborative interactions (Tomasello et al., 2005). Shared attention and joint intention may have evolved in humans as a means to communicate and cooperate with others through gaze following, and is thought to have influenced the evolution of human eye morphology to expose the white sclera around the iris (Kobayashi & Kohshima, 1997). Having a conspicuous eye that makes gaze easier to track would benefit those engaging in shared intentionality.

Unlike other corvids, jackdaws have pale irises that may facilitate the ability to track eye/head movements. von Bayern and Emery (2009a) have suggested that the pale iris may have evolved as a salient signal specifically to communicate within monogamous pairs for which successful reproduction may be dependent on coordinating actions such as finding food, nest building and defence or feeding young. In support of this proposal, jackdaws presented with an object choice task chose the correct food location only when paired with their mated partner, suggesting this task was performed cooperatively between pairs (von Bayern & Emery, 2009b). Ravens, which have dark eyes, failed a same-species object choice task (although it should be noted that ravens in monogamous pairs were not tested in a cooperative

framework as the jackdaws were; Schloegl et al., 2008a). It is unknown why some birds have evolved pale or brightly coloured irises, and no relationship has been found between breeding system and iris colour in passerine birds (Craig & Hulley, 2004), although this conclusion must remain tentative as the study did not control for phylogeny. There are also not enough comparative studies available to investigate whether sensitivity to gaze is more prominent in birds with brightly coloured eyes, or in monogamous species. One possibility is that jackdaws evolved pale irises independently of gaze following or breeding system. Therefore, rather than being a signal that evolved specifically between sender and receiver for the purpose of communication, the pale iris may be a cue (information can be extracted by the receiver) which could enhance gaze sensitivity between conspecifics. Alternatively, iris colour in jackdaws may not be related to success in gaze following tasks. It is also unclear whether the cues given by the demonstrator jackdaw in the object choice task were from the eyes, head movement or body positioning, illustrating the lack of information in the literature regarding the cues that conspecifics may or may not be using in these tasks. In fact, we argue below that animals with laterally placed eyes will have difficulty using eye movements from conspecifics for cues in gaze following (see following section).

Ultimate factors such as predation rates, individual experience, foraging behaviours, social systems and mating systems may influence proximate mechanisms including the cognitive processes by which an animal processes information obtained from gaze cues. The dynamics of social interactions may select for the evolution of cognitive mechanisms enabling more flexible, complex forms of gaze following. Studies on conspecific gaze following in various social contexts may thus enable us to examine the interaction between sociality and cognition.

Animals' responses during experiments will also often be dependent on the specific gaze cues presented (e.g. head orientation, size, colour or shape of the eyes), as demonstrated in many gaze aversion tasks (e.g. Burger et al., 1991; Carter et al., 2008; Jones, 1980; Scaife, 1976b). However, gaze following tasks often assume that the cues presented to subjects reflect those the study species uses for gaze following under natural conditions, which may not be the case. Confounding factors, such as species differences in visual configuration and hence different responses to the experimental stimuli used as gaze cues, should also be considered when interpreting results from the existing literature, and when designing gaze following experiments.

SENSORY ARCHITECTURE AND CUE INFORMATION

Consideration of sensory systems is essential to understanding instances of gaze sensitivity across taxa. For example, gaze sensitivity tasks initially designed to test underlying cognitive mechanisms in humans and other primates were designed for species with very specific visual systems: having forward-facing eyes allows gaze cues to be presented as head turning and orienting in a fixed direction, or presented as the orientation of both eyes in one direction. While there is extensive work on the gaze cues used by primates (Tomasello et al., 2007), and how the eyes have evolved as a signal in humans (Kobayashi & Kohshima, 1997, 2001), little is known about how other animals' visual systems are configured and how they respond to different cues that could be used in gaze sensitivity contexts (e.g. eye and head movements). This is particularly important as the number of species tested in gaze sensitivity tasks broadens. Existing studies include mammals with laterally placed eyes (i.e. goats, Kaminski et al., 2005; horses, *Equus caballus*, Proops & McComb, 2010), as well as reptiles (e.g. Wilkinson et al., 2010) and birds (e.g. Loretto et al., 2010; Kehmeier et al., 2011). All these species have very different visual systems. These differences are likely

to influence whether test subjects can perceive the gaze cues presented in experiments. We use birds as models to discuss the influence of visual architecture on gaze sensitivity because of the relatively large comparative literature on the avian visual system. However, when possible, we discuss the visual systems of other vertebrates. Birds show a high degree of interspecific variability in visual systems (Meyer, 1977; Martin, 2007) that is also present in other taxa (i.e. several species of birds, mammals and reptiles have laterally placed eyes, while others have frontally placed eyes). Therefore, the conclusions derived from the following discussion can be applied to other vertebrate taxa subject to gaze sensitivity studies. Our main argument is that our understanding of gaze sensitivity would benefit enormously if behavioural and cognitive studies are accompanied by a detailed characterization of the study species' visual architecture. This will determine what cues are available to indicate gaze direction and hence to what cues conspecifics or heterospecifics are sensitive.

Visual Architecture

Of the many components of the visual system, the following are likely to play a particularly relevant role in gaze sensitivity: position of the orbits, visual field configuration, degree of eye movements, and type, position and number of retinal specializations. We briefly explain each of these sensory components. Different species vary in their degree of orbit convergence (i.e. position of orbits in the skull) and thus in the extent of their binocular, lateral and blind fields around their heads (i.e. visual field configuration; Iwaniuk, Heesy, Hall, & Wylie, 2008; Martin, 2007). The placement of the orbits affects the general position of gaze in visual space as well as from where other animals can detect gaze. Bird species with more frontally placed eyes would tend to have wider binocular fields than species with more laterally placed eyes, when the eyes are at rest (Iwaniuk et al., 2008). A similar pattern has been found in mammals (Heesy, 2004). However, the degree of eye movement varies substantially between species (Fernández-Juricic, O'Rourke, & Pitlik, 2010; Martin, 2007), which can lead to variations in the visual field configuration. For example, some species can barely move their eyes (e.g. owls; Martin, 1984), whereas others with laterally placed eyes can converge and diverge their eyes (towards and away from their bills, respectively) to the point that they can have binocular fields the size of those with frontally placed eyes and extremely narrow blind areas that increase their fields of view around their heads (sparrows, Fernández-Juricic et al., 2011; Fernández-Juricic, Gall, Dolan, Tisdale, & Martin, 2008). Similar ranges in the degree of eye movement can be found in other vertebrates. For instance, chameleons can move their eyes about 180°, whereas guinea pigs can only move their eyes about 2° (Kim, 2013; Ott, 2001). These visual field configuration changes have important functional implications for enhancing food search (i.e. widening binocular fields) and predator detection (i.e. widening lateral areas), two relevant cues in gaze sensitivity scenarios.

The position of the orbits on the head also affects where potential gaze cues are available, and therefore whether other animals can perceive eye movements. For animals with frontally placed eyes, eye movements can best be perceived from the front, where both eyes can be seen (Fig. 3a). In contrast, eye movements in laterally eyed animals can best be perceived from the side, making only one eye visible from this perspective (Fig. 3a). This has important implications if an animal with laterally placed eyes is trying to detect the gaze of a conspecific that can move its eyes. If the animal is looking at the conspecific from the side, only one eye is visible. The position of the other eye is unknown to the conspecific and this can lead to ambiguity of gaze direction (Fig. 3a).

Nevertheless, the size of the visual field only describes the volume of visual space animals can perceive around their heads as a result of the projection of their retinas, but not the quality of vision. Visual performance varies in different parts of the visual field because of changes in the density of photoreceptors (i.e. involved in phototransduction) and retinal ganglion cells (i.e. involved in the transfer of information from the retina to visual centres in the brain) across the retina (Hughes, 1977). Areas of the retina with higher density of photoreceptors and retinal ganglion cells are known as retinal specializations. These retinal specializations project into a specific part of the visual field and provide higher quality information (e.g. higher visual resolution) than other parts of the retina (Collin, 1999). The retinal specializations are thought to be the centres of visual attention (Bisley, 2011). In other words, when an animal detects a visual stimulus in a sector of the visual field that is outside of the retinal specialization, it will move its head and eyes to align the retinal specialization with that object and collect high-quality information.

Retinal specializations vary in type, size, position and number (Meyer, 1977). For instance, the fovea is a retinal specialization characterized by an invagination of the retinal tissue whose centre provides the highest visual resolution (Walls, 1942). Foveae are present in many vertebrates (Duijm, 1959; Hughes, 1977; Walls, 1942) such as some primates and birds, but also in some canids and fish (Collin, Lloyd, & Wagner, 2000; Curcio et al., 1991; Dolan & Fernández-Juricic, 2010; Packer, Hendrickson, & Curcio, 1989; Peichl, 1992). The fovea projects into a smaller portion of the visual field than the visual streak, which is another retinal specialization that consists of an enlargement of the retinal tissue forming a horizontal band of high visual resolution across the central axis of the whole retina (Walls, 1942). Various vertebrate species have been found to have visual streaks (Hughes, 1977), such as horses, goats and dogfish (Bozzano, 2004; Hughes & Whitteridge, 1973; Querubin, Lee, Provis, & O'Brien, 2009). Additionally, the position and number of retinal specializations can affect the direction of gaze. For instance, some Passeriformes tend to have a single fovea projecting into the lateral field (Fernández-Juricic et al., 2011), making individuals use their lateral fields (i.e. aligning their heads laterally in relation to the object of visual interest) to explore objects visually (e.g. zebra finch, *Taeniopygia guttata*; Bischof, 1988). However, some diurnal raptors have two foveae, one central projecting to the lateral field and one temporal projecting into the binocular field (Fite & Rosenfield-Wessels, 1975; Reymond, 1985). During a chase, raptors align the fovea projecting frontally into the binocular field with the prey when close to catching it (Tucker, 2000). Thus, depending on the configuration of the visual field and the retina, the behaviours associated with gaze direction would vary between species. Variations in the number and position of the retinal specializations are also present in other vertebrates; for instance, wolves, *Canis lupus*, have a horizontal streak with a temporally placed fovea (Peichl, 1992) whereas the pigtailed macaque, *Macaca nemestrina*, has a single fovea (Packer et al., 1989).

Visual Perception in a Gaze Following Context

Two of the most important visual tasks for animals are visual search (i.e. looking for an object in visual space that is absent; such as searching for predators) and visual fixation (i.e. focusing gaze on an object that is present in visual space and gathering high-quality visual information from it with the retinal specialization, such as tracking an approaching predator). From the perspective of gaze sensitivity, visual fixation is a key process as it indicates the main centre of visual attention (Bisley, 2011). Visual fixation is associated with specific behavioural patterns (e.g. eye and head movements), which are expected to be the cues that other animals would use




Figure 3. (a) In animals with frontally placed eyes, the orientation of both eyes (as cues for gaze following) is most easily seen from a frontal view, whereas in animals with laterally placed eyes, eye orientation is more salient from the side but is partial as only one eye can be seen. (b) Visual fixation strategies proposed for bird species with laterally placed eyes. (I) locking the gaze on an object with a single fovea using the monocular field of one eye; (II) quickly alternating between the two foveae using the monocular fields of both eyes (see text for details).

during gaze detection. However, variations in the visual architecture mentioned above are likely to modify these behavioural patterns (or cues) in different ways depending on the position of the projection of the retinal specialization in visual space. Therefore, understanding visual system configuration and fixation should be two essential elements when determining the gaze cues to which animals are sensitive.

For example, humans have frontally placed orbits with a large degree of eye movement. In humans, the fovea is positioned at approximately the centre of the retina, hence projecting into the binocular field (Fig. 3a). When humans fixate, both foveae align with the object of interest with a steady gaze (Fig. 3b). When an object is static, human fixation is associated with a decrease in head movements and is fine-tuned with the eyes 'locked' on the target of attention (although the eyes still engage in very subtle movements; Martinez-Conde, 2005). A similar visual fixation strategy is present in other vertebrates such as dogs (Somppi, Tornqvist, Hanninen, Krause, & Vainio, 2012). The ocular fine-tuning in humans is facilitated by eye coloration, in which the iris surrounded by a clear sclera becomes a salient cue that facilitates gaze detection (Kobayashi & Kohshima, 1997). Overall, this visual and morphological configuration in humans reduces ambiguity in gaze direction cues.

However, in many species with laterally placed eyes (e.g. most birds, goats, horses; Fig. 3a), the type of retinal specialization, along with its projection, varies enormously between species. Additionally, their visual fixation strategies are not as well understood. Two visual fixation strategies have been proposed for birds with laterally placed eyes (Fig. 3b): (1) fixating only one fovea on a visual target using monocular vision (Maldonado, Maturana, & Varela, 1988), and (2) quickly alternating between the two foveae using the monocular fields of both eyes (Dawkins, 2002). The first

strategy is similar to human fixation in that it locks the gaze (in this case with only one eye) on the object of interest, thus reducing head movements (Fig. 3b). The second strategy actually increases head movements by having each eye check the object of interest repeatedly (Fig. 3b). Furthermore, there is evidence that fixation may also occur within the binocular field in species with laterally placed eyes when objects are very close by (Bloch, Rivaud, & Martinoya, 1984; Dawkins, 2002); however, it is not known whether this occurs by animals converging their eyes and thus projecting their retinal specialization into the binocular field. There is a major gap in comparative data as to how fixation strategies vary in vertebrates with different visual architecture, which would influence the cues other individuals use to assess gaze direction.

We can, however, make some predictions about the combination of sensory traits that could favour (or not) gaze sensitivity in species with laterally placed eyes and a single fovea. A large number of the species belonging to the most diverse avian order, Passeriformes, surveyed to date have a single fovea that is centrotemporally placed (Fernández-Juricic, 2012) and generally projects into the lateral visual field, but not far from the edge of the binocular field. These species have, however, different degrees of eye movement. If birds use eye movement as gaze direction cues as humans do, we would expect sensitivity to gaze cues to be more prevalent in species with a larger degree of eye movement (Fig. 3b), and particularly the ones in which the eye is visually salient owing to a differently coloured iris (e.g. jackdaws).

Even in species with salient (i.e. brightly coloured) eyes, there is a fundamental challenge: some bird species show coordinated eye movements whereas in others the two eyes move independently of one another (Bloch et al., 1984; Voss & Bischof, 2009). The implication is that, during fixation, the movement of one eye would

predict the movement of the other eye in some species, but not in others (Fig. 4). This uncertainty could translate into an ambiguous gaze direction cue, which may not favour gaze detection using only eye movement cues (Fig. 4). Evidence in species with laterally placed eyes supports the view that birds tend to move their heads more than their eyes when changing the direction of gaze (Gioanni, 1988). Consequently, we propose that in species with laterally placed eyes and a single fovea, species are more likely to be sensitive to head movement cues (e.g. head orientation, rate of change in head position, etc.) than eye movement cues. In those species that fixate by 'locking' their gaze to an object with a single fovea, the gaze cue is expected to be a pronounced decrease in head movement rate associated with a single head position aligned with the visual target. Conversely, in those species that fixate by using both foveae alternately, the gaze cue would be an increase in head movement rate associated with at least two main head orientations in which each eye aligns with the visual target.

Determining gaze cues (i.e. eye, head, body orientation postures that indicate where a conspecific is looking at) in bird species with a visual streak (e.g. Anseriformes) as the retinal specialization may be even more challenging. Most of the sensory issues described above apply, but additionally these species have a lower need to move their heads and eyes as the visual streak provides high visual resolution in a larger proportion of the visual field (the whole horizontal axis) than in species with foveae (Collin, 1999). We expect that species with visual streaks may be less sensitive to gaze cues, or would rely on less ambiguous cues, such as moving the head sideways to fixate the object with the retinal specialization of each eye alternately, therefore relying more on head orientation than head movement rate. Overall, we propose that visual architecture will influence not only the ability to perceive gaze cues, but also the types of cues associated with gaze direction that conspecifics and heterospecifics may use.

COGNITION IN GAZE SENSITIVITY

A species' visual system may influence the information made available to individuals in the form of gaze cues, and socioecological factors may determine whether adaptive information can be gained from attending to gaze cues (e.g. the location of food). Once it has been established that gaze cues are available to the subject and that they elicit a gaze response, we can investigate the cognitive mechanisms involved in processing gaze cue information that generate behavioural outputs.

The difficulty in interpreting the cognitive mechanisms a species is applying to gaze tasks is two-fold. First, if the sensory system of an animal is not considered, it is difficult to be certain that a negative result is due to the lack of a particular cognitive mechanism as opposed to a lack of sensitivity to a particular cue. Second, if a gaze cue is available and does cause a response, it remains difficult to disentangle whether a particular action (e.g. gaze following) is driven primarily by the stimulus (e.g. eye, head movement) or also by cognitive mechanisms that enable the subject to understand something about what the demonstrator can see. Seemingly complex behaviour may often be underpinned by relatively simple mechanisms. For example, stimulus-driven visual fixation processes in praying mantises generate complex, coordinated movements of the head, abdomen and prothorax when pinpointing the exact location of prey (Rossel, 1980; Yamawaki, Uno, Ikeda, & Toh, 2011). Similarly, the body and eye movements apparent when vertebrates redirect their visual attention in joint attention, gaze following or geometric gaze tests may also be driven by simple stimulus – response processes. One cannot ascribe the presence of gaze sensitivity to cognitive mechanisms such as perspective taking or attention attribution (see below) simply based on the complexity of behaviours observed when animals gather visual information. Instead, carefully designed experiments are essential if we are to




Figure 4. Gaze direction cues may have different degree of ambiguousness in animals with laterally placed eyes depending on whether a species has conjugate or nonconjugate eye movements. (a) Conjugate eye movements with eyes converging towards the bill. (b) Conjugate eye movements with both eyes looking to the right. (c) Nonconjugate eye movements where the left eye looks forward and the right eye is at rest towards the left side.

discriminate between alternative cognitive explanations. Often this means that authors must present alternative interpretations in the form of 'low-level' (e.g. simple behavioural responses or associative learning mechanisms) and 'high-level' mechanisms (e.g. perspective taking or attention attribution) because it is not always definitive which are driving the observable behaviours (e.g. Call et al., 1998; Povinelli & Eddy, 1996).

Alternative Interpretations

The majority of studies of the cognitive processing underlying gaze responses have employed gaze following paradigms, (but see von Bayern & Emery, 2009a; Call, Brauer, Kaminski, & Tomasello, 2003; Flombaum & Santos, 2005 for examples of cognitive tasks applying gaze aversion paradigms). Often these studies are unable to discount alternative cognitive interpretations for observed behaviour. For instance, individuals may succeed in a gaze following task by learning to associate finding food or an interesting object with seeing a particular gaze cue and then performing a gaze following behaviour. Alternatively, the subject may apply mechanisms such as shared attention or attention attribution. Attention attribution is similar to shared attention in that the subject appreciates where the demonstrator's attention is focused, but does not necessarily attend to the same object (e.g. von Bayern & Emery, 2009a).

Gaze following behaviours also raise the question of whether animals are capable of perspective taking. Perspective taking has been described as the ability to infer that others may see different things from what oneself sees (Flavell, 1974, 1977). For instance, in the geometric gaze task, a subject might take into account another individual's line of sight as being different from one's own in order to adjust its positioning around a barrier. In the literature on nonhuman gaze following, mechanisms such as shared attention, attention attribution and perspective taking are typically defined as distinct from theory of mind (the ability to reason about other individual's mental states, separate from one's own). Although theory of mind may guide gaze responses in humans, tasks in nonhuman animals cannot test for this when applying paradigms that involve behavioural cues such as eye gaze. Such tasks are unable to distinguish between responses to gaze cues themselves and responses to another individual's mental state. The most compelling evidence for perspective taking in gaze-related tasks comes from experiments that control for gaze cues or, in fact, any behavioural cue. For example, in studies of food-caching corvids, subjects have been presented with individuals that differ only in whether they had visual access to an object (i.e. food) or an event (i.e. caching; e.g. Bugnyar, 2010; Dally, Emery, & Clayton, 2006; Emery & Clayton, 2001), not in the gaze cues presented. Even so, it remains possible that demonstrators may provide subtle behavioural cues that indicate whether or not they saw food. Controlling for behavioural cues may be possible using robot models or video playback (Bird & Emery, 2008; Fernández-Juricic, Gilak, McDonald, Pithia, & Valcarcel, 2006; Woo & Rieucan, 2012; see also below).

Interpreting Negative Results

If negative results are obtained in gaze tasks, we should not always presume the absence of cognitive mechanisms in the context of gaze sensitivity. Instead, failure to perform successfully in gaze tasks may occur because the appropriate gaze cues were not available to the subject. Information on sensory systems is critical to determine whether the species is capable of attending to the demonstrators' gaze cues. If it is known that a species' visual configuration presents ambiguous gaze cues or none at all, then we should rule out mechanisms such as shared attention or

perspective taking, at least in the context of gaze following. Similarly, if the available gaze cues within a species have not been identified correctly, experimenters may be expecting to measure a behaviour that does not match the species' actual response type, given their visual architecture. For example, if both gaze cues and gaze responses within a species are very subtle (e.g. small eye movements), eye movement responses may be overlooked if head movements are the expected measure. Only once observable cues are shown to elicit measureable gaze responses can further behavioural data be collected to test for cognitive mechanisms. For example, behaviours such as turning back to face the demonstrator, presumably to confirm where they are looking (all great apes, Bräuer et al., 2005), or placing distractor objects close to the subject, but not in the demonstrator's line of sight (chimpanzees, Tomasello et al., 1999) may provide some support for shared attention. This may require the subject to attend reliably to where the demonstrator is looking, rather than stopping at the first interesting object.

With all this uncertainty, which tasks are the most informative for testing underlying cognitive mechanisms? Overall, the geometric gaze task may be a good test for complex processing in a gaze following context as it requires the subjects not only to follow the gaze of others, but also to act by adjusting their vantage point. This task also has the benefit of being ecologically relevant, as individuals may often encounter and move around barriers occluding their line of sight, or, as we have seen, may be important in species engaging in cache protection and pilfering (e.g. Bugnyar et al., 2004; Dally et al., 2006; Schloegl et al., 2007).

APPLICATIONS FOR GAZE RESEARCH

The socioecological, anatomical, sensory and cognitive features we discussed may influence the occurrence of gaze behaviours across taxa, but these factors are seldom considered together when designing and interpreting gaze tasks. To address this gap and gain a better understanding of the mechanisms underlying gaze sensitivity, we propose a new approach that consists of the following steps. Following these steps could improve our ability to interpret results, particularly in studies that show null results, while also contributing to comparative data available to gaze researchers to test how the features of an animal's visual system may be associated with the gaze cues and responses.

(1) Gaze researchers should study key components of the visual system of the study species (i.e. orbit orientation, visual field configuration, and type, position and number of retinal specializations, see <http://www.retinalmaps.com.au> for retinal topography maps) to establish the projections of the areas of acute vision into the visual field. This may be possible by studying species that are phylogenetically closely related to those with existing data available, or if limited in available study species, by collaborating with researchers that study visual systems. This will aid in making predictions regarding the degree of eye and head movement expected during visual fixation, and, where possible, to target species expected to display more pronounced gaze cues (e.g. head movement rates). (2) The behavioural mechanisms of visual fixation (e.g. head/eye orientation, movement rate, etc. when gaze is locked on an object) in the study species should be determined. This may involve observational data of the study species when presented with objects of interest in their line of sight and at different distances to identify head or eye movement associated with viewing these objects (Bossema & Burgler, 1980; Dawkins, 2002). Observational data in this context will further our understanding of how specific features of an animal's visual architecture relate to observable gaze cues. (3) It is also important to characterize the behaviours associated with visual fixation in different contexts; for

instance, are the gaze cues during food search and predator detection the same? (4) Once the gaze cues produced by the gazer are characterized, it should be established whether the cues identified in the previous step generate a gaze sensitivity response, and whether this differs depending on the socioecological context of the task (e.g. avoiding predator gaze versus the cooperative and competitive contexts when following conspecific gaze). To do so in a gaze following context, it may be beneficial to use conspecifics. This is important for those testing behaviour or cognition. If a species' visual fixation strategy differs from that in humans, the subjects may not associate human gaze in the same way they would a conspecific's gaze (but see von Bayern & Emery, 2009a where subjects were hand-raised by and had extensive interactions with humans prior to testing). Therefore, failure in a task may be measuring a lack of cue perception rather than a lack of a given cognitive mechanism. We recognize that some species are often not studied in a within-species context mostly because of logistical difficulties in manipulating gaze following cues. We suggest waiting until the appropriate gaze cue has been displayed by the demonstrator before recording subject gaze response. We also now have interesting tools at our disposal such as video playback, which has been successful for assessing same-species social preferences in rooks (Bird & Emery, 2008). Gaze cues can be manipulated by using animated video playback, which has been shown to be a successful stimulus for many species of fish, some bird species (e.g. *Lonchura punctulata*, *Gallus gallus*, *Taeniopygia guttata*) and Jacky dragons, *Amphibolurus muricatus*, (see Woo & Rieucan, 2012 for a review). Cue manipulation could also be applied using robotic animals (e.g. birds, Fernández-Juricic et al., 2006). This empirical approach can be easily adjusted to test the relative role of eye versus head movements in species with frontally and laterally placed eyes, the role of eye colour on gaze detection in birds, the relative role of different gaze following rules, etc. Alternatively, peep holes (a small hole in a wall or barrier through which the subject can look) are an effective method of determining to what subjects are attending and for how long (Bird & Emery, 2010; Grodzinski et al., 2012), and could be implemented to control what cues are observable by using different-sized peep holes exposing only the head or the eyes, or restricting species to use monocular vision only. Peep holes should be adjusted to the relative size of the species, as larger species (i.e. larger eye sizes) have higher visual acuity (Kiltie, 2000). This could be particularly relevant in studies comparing the performance of gaze sensitivity between species (e.g. territorial versus social).

Once gaze behaviours (i.e. gaze aversion, gaze following) have been established in response to characterized gaze cues, these can be applied to more complex tasks. For example, a task can be structured using the appropriate cue and a barrier to test geometric gaze. Although the gaze cue itself does not test cognitive mechanisms directly, understanding the gaze characteristics of the study species ensures that negative results are not due to the lack of cue perception.

CONCLUSION

In this review, we have proposed several socioecological, anatomical, sensory and cognitive factors that may explain the variation in gaze following or gaze aversion responses across species. We argue that it is critical to consider an animal's visual architecture as it will directly affect its ability to detect the targets of gaze. Gaze cues can differ between contexts within the same species, for instance whether the visual fixation strategy used by a conspecific is being presented as a cue during food search or as a cue during predator scanning. Furthermore, the gaze cues detectable between conspecifics may be different from gaze cues

presented by heterospecifics or predators. Therefore it is crucial to ensure that appropriate cues are chosen to match the context of the task. This presents researchers with a unique opportunity to test how variations in sensory systems can affect the occurrence of gaze sensitivity across species. Finally, establishing the gaze cues to which each species attends, and under what conditions, will provide robust experimental designs for gaze tasks testing cognitive mechanisms.

Acknowledgments

We are grateful to Ljerka Ostojic and Lucy Cheke for comments and discussion. Four anonymous referees provided constructive criticism and useful suggestions. This work was funded by the Zoology Balfour Fund (G.D.), The BBSRC David Phillips Research Fellowship (A.T.) and the National Science Foundation (E.F.J., S.B.).

References

- Amici, F., Aureli, F., Visalberghi, E., & Call, J. (2009). Spider monkeys (*Ateles geoffroyi*) and capuchin monkeys (*Cebus apella*) follow gaze around barriers: evidence for perspective taking? *Journal of Comparative Psychology*, *123*, 368–374.
- Anderson, J. R., Montant, M., & Schmitt, D. (1996). Rhesus monkeys fail to use gaze direction as an experimenter-given cue in an object-choice task. *Behavioural Processes*, *37*, 47–55.
- Anderson, J. R., Sallaberry, P., & Barbier, H. (1995). Use of experimenter-given cues during object-choice tasks by capuchin monkeys. *Animal Behaviour*, *49*, 201–208.
- Baron-Cohen, S. (1994). How to build a baby that can read minds: cognitive mechanisms in mindreading. *Cahiers de Psychologie Cognitive*, *13*, 513–552.
- von Bayern, A. M., & Emery, N. J. (2009a). Jackdaws respond to human attentional states and communicative cues in different contexts. *Current Biology*, *19*, 602–606.
- von Bayern, A. M., & Emery, N. J. (2009b). Bonding, mentalising and rationality. In S. Watanabe (Ed.), *Irrational humans, rational animals* (pp. 287–303). Tokyo: Keio University Press.
- Bird, C. D., & Emery, N. J. (2008). Using video playback to investigate the social preferences of rooks, *Corvus frugilegus*. *Animal Behaviour*, *76*, 679–687.
- Bird, C. D., & Emery, N. J. (2010). Rooks perceive support relations similar to six-month-old babies. *Proceedings of the Royal Society B*, *277*, 147–151.
- Bischof, H. J. (1988). The visual field and visually guided behavior in the zebra finch (*Taeniopygia guttata*). *Journal of Comparative Physiology*, *163*, 329–337.
- Bisley, J. W. (2011). The neural basis of visual attention. *Journal of Physiology*, *589*, 49–57.
- Bloch, S., Rivaud, S., & Martinoya, C. (1984). Comparing frontal and lateral viewing in the pigeon. III. Different patterns of eye movements for binocular and monocular fixation. *Behavioural Brain Research*, *13*, 173–182.
- Bock, S. W., Dicke, P., & Thier, P. (2008). How precise is gaze following in humans? *Vision Research*, *48*, 946–957.
- Bossema, I., & Burgler, R. R. (1980). Communication during monocular and binocular looking in European jays (*Garrulus g. glandarius*). *Behaviour*, *74*, 274–283.
- Bozzano, A. (2004). Retinal specialisations in the dogfish *Centroscyllium coelepis* from the Mediterranean deep-sea. *Scientia Marina*, *68*, 185–195.
- Bräuer, J., Call, J., & Tomasello, M. (2004). Visual perspective taking in dogs (*Canis familiaris*) in the presence of barriers. *Applied Animal Behaviour Science*, *88*, 299–317.
- Bräuer, J., Call, J., & Tomasello, M. (2005). All great ape species follow gaze to distant locations and around barriers. *Journal of Comparative Psychology*, *119*, 145–154.
- Bugnyar, T. (2010). Knower-guesser differentiation in ravens: others' viewpoints matter. *Proceedings of the Royal Society B*, *278*, 634–640.
- Bugnyar, T., & Kotrschal, K. (2002). Observational learning and the raiding of food caches in ravens, *Corvus corax*: is it 'tactical' deception? *Animal Behaviour*, *64*, 185–195.
- Bugnyar, T., Stowe, M., & Heinrich, B. (2004). Ravens, *Corvus corax*, follow gaze direction of humans around obstacles. *Proceedings of the Royal Society B*, *271*, 1331–1336.
- Burger, J., & Gochfeld, M. (1993). The importance of the human face in risk perception by black iguanas, *Ctenosaura similis*. *Journal of Herpetology*, *27*, 426–430.
- Burger, J., Gochfeld, M., & Murray, B. G., Jr. (1991). Role of a predator's eye size in risk perception by basking black iguana, *Ctenosaura similis*. *Animal Behaviour*, *42*, 471–476.
- Burkart, J. M., & Heschl, A. (2007). Understanding visual access in common marmosets, *Callithrix jacchus*: perspective taking or behaviour reading? *Animal Behaviour*, *73*, 457–469.
- Call, J., Agnetta, B., & Tomasello, M. (2000). Cues that chimpanzees do and do not use to find hidden objects. *Animal Cognition*, *3*, 23–34.

- Call, J., Brauer, J., Kaminski, J., & Tomasello, M. (2003). Domestic dogs (*Canis familiaris*) are sensitive to the attentional state of humans. *Journal of Comparative Psychology*, *117*, 257–263.
- Call, J., Hare, B. A., & Tomasello, M. (1998). Chimpanzee gaze following in an object-choice task. *Animal Cognition*, *1*, 89–99.
- Carter, J., Lyons, N. J., Cole, H. L., & Goldsmith, A. R. (2008). Subtle cues of predation risk: starlings respond to a predator's direction of eye-gaze. *Proceedings of the Royal Society B*, *275*, 1709–1715.
- Collin, S. P. (1999). Behavioural ecology and retinal cell topography. In S. Archer, M. Djamgoz, E. Loew, J. C. Partridge, & S. Vallerga (Eds.), *Adaptive mechanisms in the ecology of vision* (pp. 509–535). Dordrecht: Kluwer Academic.
- Collin, S. P., Lloyd, D. J., & Wagner, H. J. (2000). Foveate vision in deep-sea teleosts: a comparison of primary visual and olfactory inputs. *Philosophical Transactions of the Royal Society B*, *355*, 1315–1320.
- Coss, R. G. (1979). Delayed plasticity of an instinct: recognition and avoidance of 2 facing eyes by the jewel fish. *Developmental Psychobiology*, *12*, 335–345.
- Craig, A. J. F. K., & Hulley, P. E. (2004). Iris colour in passerine birds: why be bright-eyed? *South African Journal of Science*, *100*, 584–588.
- Curcio, C. A., Allen, K. A., Sloan, K. R., Lerea, C. L., Hurley, J. B., Klock, I. B., et al. (1991). Distribution and morphology of human cone photoreceptors stained with anti-blue opsin. *Journal of Comparative Neurology*, *312*, 610–624.
- Dally, J. M., Emery, N. J., & Clayton, N. S. (2005). Cache protection strategies by western scrub-jays, *Aphelocoma californica*: implications for social cognition. *Animal Behaviour*, *70*, 1251–1263.
- Dally, J. M., Emery, N. J., & Clayton, N. S. (2006). Food-caching western scrub-jays keep track of who was watching when. *Science*, *312*, 1662–1665.
- Dawkins, M. (2002). What are birds looking at? Head movements and eye use in chickens. *Animal Behaviour*, *63*, 991–998.
- Dolan, T., & Fernández-Juricic, E. (2010). Retinal ganglion cell topography of five species of ground-foraging birds. *Brain, Behavior and Evolution*, *75*, 111–121.
- Duijm, M. (1959). On the position of a ribbon-like central area in the eye of some birds. *Archives Néerlandaises de Zoologie*, *13*, 145.
- Emery, N. J. (2000). The eyes have it: the neuroethology, function and evolution of social gaze. *Neurosciences and Biobehavioral Reviews*, *24*, 581–604.
- Emery, N. J., & Clayton, N. S. (2001). Effects of experience and social context on prospective caching strategies by scrub jays. *Nature*, *414*, 443–446.
- Emery, N. J., Lorincz, E. N., Perrett, D. I., Oram, M. W., & Baker, C. I. (1997). Gaze following and joint attention in rhesus monkeys (*Macaca mulatta*). *Journal of Comparative Psychology*, *111*, 286–293.
- Fernández-Juricic, E. (2012). Sensory basis of vigilance behavior in birds: synthesis and future prospects. *Behavioural Processes*, *89*, 143–152.
- Fernández-Juricic, E., Erichsen, J. T., & Kacelnik, A. (2004). Visual perception and social foraging in birds. *Trends in Ecology & Evolution*, *19*, 25–31.
- Fernández-Juricic, E., Gall, M. D., Dolan, T., O'Rourke, C. T., Thomas, S., & Lynch, J. R. (2011). Visual systems and vigilance behaviour of two ground-foraging avian prey species: white-crowned sparrows and California towhees. *Animal Behaviour*, *81*, 705–713.
- Fernández-Juricic, E., Gall, M. D., Dolan, T., Tisdale, V., & Martin, G. (2008). The visual fields of two ground-foraging birds, house finches and house sparrows, allow for simultaneous foraging and anti-predator vigilance. *Ibis*, *150*, 779–787.
- Fernández-Juricic, E., Gilak, N., Mcdonald, J. C., Pithia, P., & Valcarcel, A. (2006). A dynamic method to study the transmission of social foraging information in flocks using robots. *Animal Behaviour*, *71*, 901–911.
- Fernández-Juricic, E., & Kowalski, V. (2011). Where does a flock end from an information perspective? A comparative experiment with live and robotic birds. *Behavioral Ecology*, *22*, 1304–1311.
- Fernández-Juricic, E., O'Rourke, C., & Pitlik, T. (2010). Visual coverage and scanning behavior in two corvid species: American crow and Western scrub jay. *Journal of Comparative Physiology A*, *196*, 879–888.
- Fernández-Juricic, E., Smith, R., & Kacelnik, A. (2005). Increasing the costs of conspecific scanning in socially foraging starlings affects vigilance and foraging behaviour. *Animal Behaviour*, *69*, 73–81.
- Fitch, W. T., Huber, L., & Bugnyar, T. (2010). Social cognition and the evolution of language: constructing cognitive phylogenies. *Neuron*, *65*, 795–814.
- Fite, K. V., & Rosenfield-Wessels, S. (1975). A comparative study of deep avian foveas. *Brain, Behavior and Evolution*, *12*, 97–115.
- Flavell, J. (1974). The development of inferences about others. In T. Mischel (Ed.), *Understanding other persons* (pp. 66–116). Oxford: Blackwell.
- Flavell, J. (1977). The development of knowledge about visual perception. *The Nebraska Symposium on Motivation*, *25*, 43–76.
- Flombaum, J. I., & Santos, L. R. (2005). Rhesus monkeys attribute perceptions to others. *Current Biology*, *15*, 447–452.
- Gioanni, H. (1988). Stabilizing gaze reflexes in the pigeon (*Columba livia*). I. Horizontal and vertical optokinetic eye (OKN) and head (OCR) reflexes. *Experimental Brain Research*, *69*, 567–582.
- Grodzinski, U., Watanabe, A., & Clayton, N. S. (2012). Peep to pilfer: what scrub-jays like to watch when observing others. *Animal Behaviour*, *83*, 1253–1260.
- Hampton, R. R. (1994). Sensitivity to information specifying the line of gaze of humans in sparrows (*Passer domesticus*). *Behaviour*, *130*, 41–51.
- Hare, B., Call, J., Agnetta, B., & Tomasello, M. (2000). Chimpanzees know what conspecifics do and do not see. *Animal Behaviour*, *59*, 771–785.
- Hare, B., Call, J., & Tomasello, M. (2006). Chimpanzees deceive a human competitor by hiding. *Cognition*, *101*, 495–514.
- Hare, B., & Tomasello, M. (2004). Chimpanzees are more skillful in competitive than in cooperative cognitive tasks. *Animal Behaviour*, *68*, 571–581.
- Hauser, M. D. (1992). Costs of deception: cheaters are punished in rhesus monkeys (*Macaca mulatta*). *Proceedings of the National Academy of Sciences of the United States of America*, *89*, 12137–12139.
- Hauser, M. D., Teixidor, P., Fields, L., & Flaherty, R. (1993). Food-elicited calls in chimpanzees: effects of food quantity and divisibility. *Animal Behaviour*, *45*, 817–819.
- Heesy, C. P. (2004). On the relationship between orbit orientation and binocular visual field overlap in mammals. *The Anatomical Record Part A: Discoveries in Molecular, Cellular, and Evolutionary Biology*, *281*, 1104–1110.
- Hughes, A. (1977). The topography of vision in mammals of contrastive life style: comparative optics and retinal organisation. In F. Crescitelli (Ed.), *The visual system in vertebrates* (pp. 615–756). New York: Springer-Verlag.
- Hughes, A., & Whitteridge, D. (1973). The receptive fields and topographical organization of goat retinal ganglion cells. *Vision Research*, *13*, 1101–1114.
- Iwaniuk, A. N., Heesy, C. P., Hall, M. I., & Wylie, D. R. (2008). Relative Wulst volume is correlated with orbit orientation and binocular visual field in birds. *Journal of Comparative Physiology A*, *194*, 267–282.
- Jones, R. B. (1980). Reactions of male domestic chicks to two-dimensional eye-like shapes. *Animal Behaviour*, *28*, 212–218.
- Kaminski, J., Riedel, J., Call, J., & Tomasello, M. (2005). Domestic goats, *Capra hircus*, follow gaze direction and use social cues in an object choice task. *Animal Behaviour*, *69*, 11–18.
- Kehmeier, S., Schloegl, C., Scheiber, I., & Weiß, B. (2011). Early development of gaze following into distant space in juvenile greylag geese (*Anser anser*). *Animal Cognition*, *14*, 477–485.
- Kiltie, R. A. (2000). Scaling of visual acuity with body size in mammals and birds. *Functional Ecology*, *14*, 226–234.
- Kim, J. (2013). Tonic eye movements induced by bilateral and unilateral galvanic vestibular stimulation (GVS) in guinea pigs. *Brain Research Bulletin*, *90*, 72–78.
- Kobayashi, H., & Kohshima, S. (1997). Unique morphology of the human eye. *Nature*, *387*, 767–768.
- Kobayashi, H., & Kohshima, S. (2001). Unique morphology of the human eye and its adaptive meaning: comparative studies on external morphology of the primate eye. *Journal of Human Evolution*, *40*, 419–435.
- Liebal, K., & Kaminski, J. (2012). Gibbons (*Hylobates pileatus*, *H. moloch*, *H. lar*, *Symphalangus syndactylus*) follow human gaze, but do not take the visual perspective of others. *Animal Cognition*, *15*, 1211–1216.
- Lima, S. L. (1987). Vigilance while feeding and its relation to the risk of predation. *Journal of Theoretical Biology*, *124*, 303–316.
- Loretto, M. C., Schloegl, C., & Bugnyar, T. (2010). Northern bald ibises follow others' gaze into distant space but not behind barriers. *Biology Letters*, *6*, 14–17.
- Maldonado, P. E., Maturana, H., & Varela, F. J. (1988). Frontal and lateral visual system in birds. Frontal and lateral gaze. *Brain, Behavior and Evolution*, *32*, 57–62.
- Martin, G. R. (1984). The visual fields of the tawny owl, *Strix aluco* L. *Vision Research*, *24*, 1739–1751.
- Martin, G. (2007). Visual fields and their functions in birds. *Journal of Ornithology*, *148*, 547–562.
- Martinez-Conde, S. (2005). The role of eye movement during visual fixation. *Barrow Quarterly*, *21*, 44–48.
- Meyer, D. B. C. (1977). The avian eye and its adaptations. In F. Crescitelli (Ed.), *The visual system in vertebrates* (pp. 549–612). New York: Springer.
- Mulcahy, N. J., & Hedge, V. (2012). Are great apes tested with an object-object-choice task? *Animal Behaviour*, *83*, 313–321.
- Ott, M. (2001). Chameleons have independent eye movements but synchronise both eyes during saccadic prey tracking. *Experimental Brain Research*, *139*, 173–179.
- Packer, O., Hendrickson, A. E., & Curcio, C. A. (1989). Photoreceptor topography of the retina in the adult pigtail macaque (*Macaca nemestrina*). *Journal of Comparative Neurology*, *288*, 165–183.
- Peichl, L. (1992). Topography of ganglion cells in the dog and wolf retina. *Journal of Comparative Neurology*, *324*, 603–620.
- Povinelli, D. J., & Eddy, T. J. (1996). Chimpanzees: joint visual attention. *Psychological Science*, *7*, 129–135.
- Proops, L., & McComb, K. (2010). Attributing attention: the use of human-given cues by domestic horses (*Equus caballus*). *Animal Cognition*, *13*, 197–205.
- Querubin, A., Lee, H. R., Provis, J. M., & O'Brien, K. M. (2009). Photoreceptor and ganglion cell topographies correlate with information convergence and high acuity regions in the adult pigeon (*Columba livia*) retina. *Journal of Comparative Neurology*, *517*, 711–722.
- Reymond, L. (1985). Spatial visual acuity of the eagle *Aquila audax*: a behavioural, optical and anatomical investigation. *Vision Research*, *25*, 1477–1491.
- Rossel, S. (1980). Foveal fixation and tracking in the praying mantis. *Journal of Comparative Physiology*, *139*, 307–331.
- le Roux, R. A., Snyder-Mackler, N., Roberts, E. K., Beehner, J. C., & Bergman, T. J. (2013). Evidence for tactical concealment in a wild primate. *Nature Communications*, *4*, 1462.
- Scaife, M. (1976a). The response to eye-like shapes by birds. I. The effect of context: a predator and a strange bird. *Animal Behaviour*, *24*, 195–199.
- Scaife, M. (1976b). The response to eye-like shapes by birds II. The importance of staring, pairedness and shape. *Animal Behaviour*, *24*, 200–206.
- Schloegl, C., Kotrschal, K., & Bugnyar, T. (2007). Gaze following in common ravens, *Corvus corax*: ontogeny and habituation. *Animal Behaviour*, *74*, 769–778.
- Schloegl, C., Kotrschal, K., & Bugnyar, T. (2008a). Do common ravens (*Corvus corax*) rely on human or conspecific gaze cues to detect hidden food? *Animal Cognition*, *11*, 231–241.

- Schloegl, C., Kotrschal, K., & Bugnyar, T. (2008b). Modifying the object-choice task: is the way you look important for ravens? *Behavioural Processes*, *77*, 61–65.
- Schmidt, J., Scheid, C., Kotrschal, K., Bugnyar, T., & Schloegl, C. (2011). Gaze direction – a cue for hidden food in rooks (*Corvus frugilegus*)? *Behavioural Processes*, *88*, 88–93.
- Shaw, R. C., & Clayton, N. S. (2012). Eurasian jays, *Garrulus glandarius*, flexibly switch caching and pilfering tactics in response to social context. *Animal Behaviour*, *84*, 1191–1200.
- Shaw, R. C., & Clayton, N. S. (2013). Careful cachers and prying pilferers: Eurasian jays (*Garrulus glandarius*) limit auditory information available to competitors. *Proceedings of the Royal Society B*, *280*, 20122238.
- Sommer, V., & Reichard, U. (2000). Rethinking monogamy: the gibbon case. In P. M. Kappeler (Ed.), *Primate males: Causes and consequences of variation in group composition* (pp. 159–168). Cambridge: Cambridge University Press.
- Somppi, S., Tornqvist, H., Hanninen, L., Krause, C., & Vainio, O. (2012). Dogs do look at images: eye tracking in canine cognition research. *Animal Cognition*, *15*, 163–174.
- Stevens, M., Hopkins, E., Hinde, W., Adcock, A., Connolly, Y., Troscianko, T., et al. (2007). Field experiments on the effectiveness of 'eyespot' as predator deterrents. *Animal Behaviour*, *74*, 1215–1227.
- Stulp, G., Emery, N. J., Verhulst, S., & Clayton, N. S. (2009). Western scrub-jays conceal auditory information when competitors can hear but cannot see. *Biology Letters*, *5*, 583–585.
- Teufel, C., Alexis, D. M., Clayton, N. S., & Davis, G. (2010). Mental-state attribution drives rapid, reflexive gaze following. *Attention, Perception Psychophysics*, *72*, 695–705.
- Thornton, A., & McAuliffe, K. (2012). Teaching can teach us a lot. *Animal Behaviour*, *83*, e6–e9.
- Thornton, A., & Raihani, N. J. (2008). The evolution of teaching. *Animal Behaviour*, *75*, 1823–1836.
- Tomasello, M., Call, J., & Hare, B. (1998). Five primate species follow the visual gaze of conspecifics. *Animal Behaviour*, *55*, 1063–1069.
- Tomasello, M., & Carpenter, M. (2005). The emergence of social cognition in three young chimpanzees. *Monographs of the Society for Research in Child Development*, *70*, 133–152.
- Tomasello, M., Carpenter, M., Call, J., Behne, T., & Moll, H. (2005). Understanding and sharing intentions: the origins of cultural cognition. *Behavioral and Brain Sciences*, *28*, 675–691.
- Tomasello, M., Hare, B., & Agnetta, B. (1999). Chimpanzees, *Pan troglodytes*, follow gaze direction geometrically. *Animal Behaviour*, *58*, 769–777.
- Tomasello, M., Hare, B., Lehmann, H., & Call, J. (2007). Reliance on head versus eyes in the gaze following of great apes and human infants: the cooperative eye hypothesis. *Journal of Human Evolution*, *52*, 314–320.
- Tomonaga, M., & Imura, T. (2010). Visual search for human gaze direction by a chimpanzee (*Pan troglodytes*). *PLoS One*, *5*, e9131.
- Tucker, V. A. (2000). The deep fovea, sideways vision and spiral flight paths in raptors. *Journal of Experimental Biology*, *203*, 3745–3754.
- Voss, J., & Bischof, H. J. (2009). Eye movements of laterally eyed birds are not independent. *Journal of Experimental Biology*, *212*, 1568–1575.
- Walls, L. (1942). *The vertebrate eye and its adaptive radiation*. Cranbrook Institute of Science. Bulletin no. 19.
- Watve, M., Thakar, J., Kale, A., Puntambekar, S., Shaikh, I., Vaze, K., et al. (2002). Bee-eaters (*Merops orientalis*) respond to what a predator can see. *Animal Cognition*, *5*, 253–259.
- Wilkinson, A., Mandl, I., Bugnyar, T., & Huber, L. (2010). Gaze following in the red-footed tortoise (*Geochelone carbonaria*). *Animal Cognition*, *13*, 765–769.
- Woo, K. L., & Rieucau, G. (2012). From dummies to animations: a review of computer-animated stimuli used in animal behavior studies. *Behavioral Ecology and Sociobiology*, *65*, 1671–1685.
- Yamawaki, Y., Uno, K., Ikeda, R., & Toh, Y. (2011). Coordinated movements of the head and body during orienting behaviour in the praying mantis *Tenodera arifolia*. *Journal of Insect Physiology*, *57*, 1010–1016.